

Alzheimer's disease via enhanced calcium signaling caused by the decrease of endoplasmic reticulum–mitochondrial distance



Hong Qi ^{a,*}, Jianwei Shuai ^{b,c,*}

^aComplex Systems Research Center, Shanxi University, Taiyuan 030006, PR China

^bDepartment of Physics, Xiamen University, Xiamen 361005, PR China

^cState Key Laboratory of Cellular Stress Biology, Innovation Center for Cell Signaling Network, Xiamen University, Xiamen 361005, PR China

ARTICLE INFO

Article history:

Received 9 November 2015

Accepted 30 January 2016

ABSTRACT

It has long been recognized that Ca^{2+} dysregulation is relevant to the initiation of Alzheimer's disease (AD), and most recent works have suggested that increased cross-talk between endoplasmic reticulum (ER) and mitochondria plays an important role in the pathogenesis of the disease. However, the detailed mechanism involved has not been fully elucidated. Owing to its importance in the regulation of Ca^{2+} signaling, ER-mitochondrial distance in the neurons is tightly controlled in the physiological conditions. When the distance is decreased, Ca^{2+} overload occurs both in the cytosol and mitochondria. The cytosolic Ca^{2+} overload can (1) hyperactivate Ca^{2+} -dependent enzymes, which in turn regulate activities of pro-apoptotic BCL-2 family proteins, causing mitochondrial outer membrane permeabilization and thereby resulting in the release of cytochrome c to activate caspase-3; (2) indirectly activate caspase-3 through the activation of caspase-12; and (3) promote the production and aggregation of β -amyloid. The three pathways eventually trigger neuronal apoptotic cell death. The mitochondrial Ca^{2+} overload can lead to increased generation of reactive oxygen species, inducing the opening of the mitochondrial permeability transition pore and ultimately causing neuronal apoptotic and necrotic cell death. The resultant death of neurons which are responsible for memory and cognition would contribute to the pathogenesis of AD. Therefore, we propose that the reduction in the distance between ER and mitochondria may be implicated in AD pathology by enhanced Ca^{2+} signaling, which provides a more complete picture of the Ca^{2+} hypothesis of AD.

© 2016 Elsevier Ltd. All rights reserved.

Introduction

Alzheimer's disease (AD) is the most common human neurodegenerative disorder, and is characterized by the accumulation of β -amyloid and the progressive impairment of memory and cognition that is strongly correlated with death of neurons in the hippocampus and neocortex [1–5]. Neuronal cell death may occur through apoptosis and necrosis [6,7], the former being the predominant form in AD [3,8]. Despite extensive research into the etiologies of AD, the underlying pathogenic mechanisms involved remain obscure and no scientific explanation has gained total acceptance [9]. The main hypotheses that have been put forward to explain the causes of AD include amyloid cascade hypothesis [10], endoplasmic reticulum (ER) stress hypothesis [11,12], mitochondrial cascade hypothesis [13,14], and Ca^{2+} hypothesis of AD [15–17].

* Address: Department of Physics, Xiamen University, Xiamen 361005, PR China.
Tel.: +86 139 5928 7814 (J. Shuai). Tel.: +86 0351 7019655 (H. Qi).

E-mail addresses: hongqi@sxu.edu.cn (H. Qi), jianweishuai@xmu.edu.cn (J. Shuai).

A proper increase of cytosolic Ca^{2+} concentration plays a pivotal role in regulating many neuronal functions, such as information processing, learning and memory [17]. The progressive decline in cognition is intimately linked to neuronal death that can be driven by the marked and sustained elevation of Ca^{2+} signaling [18], which forms the basis of the Ca^{2+} hypothesis of AD [19,20].

Neuronal Ca^{2+} homeostasis depends critically on the Ca^{2+} transfer from ER to mitochondria, which is ascribed to the close connection between both organelles [21,22]. Such a connection is maintained by several tethering proteins such as presenilin 2 (PS2) [23], phosphofuran acidic cluster sorting protein-2 (PACS2) [24], and mitofusin-2 (MFN2) [25]. It was revealed experimentally that a proper spacing between ER and mitochondria plays a regulatory role in Ca^{2+} signaling and that tightening of the gap facilitates mitochondrial Ca^{2+} overload and commits the cells to a cell-death pathway [26]. Most recently we demonstrated that there is an optimal distance between ER and mitochondria for the physiological Ca^{2+} signals and proposed that apoptosis induced by elevated cytosolic Ca^{2+} may be observed when the optimal distance is disturbed [27].

Indeed, Area-Gomez et al. found that ER-mitochondrial communication is increased significantly in AD [28]. Based on this finding, they proposed that AD is a disorder of ER-mitochondrial communication [29]. It is compelling that the communication is involved in the transfer of Ca^{2+} between the two organelles to carry out the regulation of Ca^{2+} signaling. This raises the question of whether AD might result from changes in ER-mitochondrial distance.

Hypothesis

The reduction in the ER-mitochondrial distance in neurons, which may be attributable to different causes (such as abnormal proteins expression, cellular stress, and aging), would result in excessive rises of Ca^{2+} within cytosol and mitochondria, which in turn activate their downstream cell death signals, contributing to the pathogenesis of AD.

Facts and evidences

Neurons have an elaborate ER network that extends throughout the cell [30]. A large portion of this network comes into close contact with mitochondria [24]. A huge body of literatures has addressed the dysfunction of ER, mitochondria and Ca^{2+} in AD. In the following, we list the facts and evidences to support our hypothesis (Fig. 1).

- (1) A correct distance between ER and mitochondria is essential for their mutual interactions, which modulate key aspects of cell physiology [25]. However, reduction in the distance

between them may occur in some pathophysiological conditions. First, ER-mitochondrial distance can be decreased by the abnormal expressions of tethering proteins: up-regulation of PS2 [23] or PACS2 [24], down-regulation of MFN2 [25]. Second, tight ER-mitochondrial coupling appears during early phases of ER stress [31]. Third, neuronal shrinkage [32,33] and increased mitochondrial size [34,35], which have been well-documented in AD brain [36,37] and are two concomitants of brain aging, may cause compact ER-mitochondrial juxtaposition, indicating that aging is the major risk factor for the development of AD [38].

- (2) If ER-mitochondrial distance is shorter than the optimal value, the amount of Ca^{2+} ions released from the ER will increase, which in turn leads to elevated Ca^{2+} in both cytosol and mitochondria [27]. Although this conclusion was obtained by ignoring the contribution of extracellular sources of Ca^{2+} in theory, such transients have been observed experimentally in neurons [39]. In addition, changes in Ca^{2+} dynamics and levels are important features of AD pathology, including enhanced Ca^{2+} release from ER [40,41], exaggerated cytosolic Ca^{2+} [42–44] and overload mitochondrial Ca^{2+} [45,46].
- (3) When cytoplasmic Ca^{2+} concentration ($[\text{Ca}^{2+}]_{\text{cyt}}$) exceeds the normal physiological level, apoptotic cell death can occur through three pathways. Firstly, elevated $[\text{Ca}^{2+}]_{\text{cyt}}$ results in the activation of caspase-12 [47], which subsequently activates caspase-3 [48,49]. This pathway is important in ER stress-induced neuronal apoptosis [49]. Secondly, two Ca^{2+} -dependent proteases, calpain [50,51] and calcineurin [52,53], are hyperactivated under pathological conditions




Fig. 1. Decreased ER-mitochondrial distance hypothesis of AD pathogenesis. ER, endoplasmic reticulum; Mt, mitochondria; $[\text{Ca}^{2+}]_{\text{cyt}}$, cytosolic Ca^{2+} concentration; $[\text{Ca}^{2+}]_{\text{Mt}}$, mitochondrial Ca^{2+} concentration; MOMP, mitochondrial outer membrane permeabilization; mPTP, mitochondrial permeability transition pore.

that involve sustained cytoplasmic Ca^{2+} overload, which in turn activates BH3-only family members Bid [54] and Bad [55], respectively. Either directly or indirectly, these two members can induce the activation of pro-apoptotic effector Bak [56,57], thereby enabling them to form pores that cause mitochondrial outer membrane permeabilization (MOMP) [58]. As a result of MOMP, cytochrome *c* is released from the mitochondrial intermembrane space into the cytosol, where cytochrome *c* promotes caspase-3 activation [56,57]. Caspase-3 cleaves a series of cellular substrates and orchestrates apoptosis [56,59]. Experimental evidences support the involvement of calpain [60] and calcineurin [53] as well as Bcl-2 family members [61] and caspase [62] in the neuronal apoptotic cell death that occurs in AD. Thirdly, high levels of $[\text{Ca}^{2+}]_{\text{cyt}}$ induces apoptosis [63] by accelerating the generation [40] and aggregation [63] of β -amyloid.

(4) Mitochondrial Ca^{2+} overload has been associated with almost all forms of cell death [64], including apoptosis and necrosis [21]. Ca^{2+} transfer from the ER to mitochondria is required to maintain normal cell bioenergetics for survival [65]. However, excessive Ca^{2+} accumulation within mitochondria can enhance the generation of reactive oxygen species [66], which form a coherent feed-forward loop with an AND input function [67], controlling the opening of mitochondrial permeability transition pore (mPTP) [66] and ultimately leading to cell death either by apoptosis or necrosis [68,69]. In other words, mPTP is capable of translating mitochondrial Ca^{2+} into a death signal only if mitochondrial Ca^{2+} overloads and results in ROS over-production. There is evidence that the absence of cyclophilin D (an integral part of mPTP) protects neurons from cell death and ameliorates learning and memory in AD [70].

(5) Both apoptosis and necrosis can significantly contribute to massive neuronal cell loss, which is a cardinal feature of AD. Apoptosis has not only been documented in studies of animal and cell-culture models of AD [1], but also of post-mortem brain tissue in AD [7]. Although many neurons may die by apoptosis in AD [3], other modes of cell death, such as necrosis, are relevant for the ultimate progression of the disease [71,72]. It has been proved that brain regions involved in learning and memory processes, including hippocampus and neocortex, are reduced in size in AD patients as the result of neuronal demise [3].

Conclusions and future perspectives

Emerging evidence suggests that AD-linked PS2 mutation also modulates the physical interaction between ER and mitochondria [23] and the increased cross-talk between the two organelles may play an important role in AD pathology [24,28]. Furthermore, considering the profound impact of ER-mitochondrial distance on Ca^{2+} signaling as well as Ca^{2+} -mediated cell death on AD pathology, it is logical to hypothesize that the decreased distance may sensitize neurons to Ca^{2+} -induced cell death, and therefore contribute to the pathogenesis of AD.

Our hypothesis can be tested by measuring and comparing the distance between ER and mitochondria, the Ca^{2+} levels within the cytosol and mitochondria, and the activities of cell death-related protein in healthy humans and AD patients. Overall, our hypothesis provides deeper insights into how Ca^{2+} dysregulation plays an early proximal and central role in AD [40,73] and clarifies a possible role of ER-mitochondrial distance in AD pathogenesis, which may suggest directions for treating the disease at an early stage.

Conflict of interest

We declare that we do not have any commercial or associative interest that represents a conflict of interest in connection with the work submitted.

Acknowledgments

This work is supported by National Natural Science Foundation of China Grants 11504214 and 31370830.

References

- [1] Mattson MP. Apoptosis in neurodegenerative disorders. *Nat Rev Mol Cell Biol* 2000;1:120–30.
- [2] Shimohama S. Apoptosis in Alzheimer's disease—an update. *Apoptosis* 2000;5:9–16.
- [3] Mattson MP. Pathways towards and away from Alzheimer's disease. *Nature* 2004;430:631–9.
- [4] Choi SH, Kim YH, Heisch M, et al. A three-dimensional human neural cell culture model of Alzheimer's disease. *Nature* 2014;515:274–8.
- [5] Lu T, Aron L, Zullo J, et al. REST and stress resistance in ageing and Alzheimer's disease. *Nature* 2014;507:448–54.
- [6] Yuan J, Lipinski M, Degterev A. Diversity in the mechanisms of neuronal cell death. *Neuron* 2003;40:401–13.
- [7] Gorman AM. Neuronal cell death in neurodegenerative diseases: recurring themes around protein handling. *J Cell Mol Med* 2008;12:2263–80.
- [8] Friedlander RM. Apoptosis and caspases in neurodegenerative diseases. *N Engl J Med* 2003;348:1365–75.
- [9] Grases F, Costa-Bauzá A, Prieto RM. A potential role for crystallization inhibitors in treatment of Alzheimer's disease. *Med Hypotheses* 2010;74:118–9.
- [10] Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 2002;297:353–6.
- [11] Lindholm D, Wootz H, Korhonen L. ER stress and neurodegenerative diseases. *Cell Death Differ* 2006;13:385–92.
- [12] Roussel BD, Krappa AJ, Miranda E, Crowther DC, Lomas DA, Marciniak SJ. Endoplasmic reticulum dysfunction in neurological disease. *Lancet Neurol* 2013;12:105–18.
- [13] Swerdlow RH, Khan SM. A "mitochondrial cascade hypothesis" for sporadic Alzheimer's disease. *Med Hypotheses* 2004;63:8–20.
- [14] Swerdlow RH, Burns JM, Khan SM. The Alzheimer's disease mitochondrial cascade hypothesis: progress and perspectives. *Biochim Biophys Acta* 2014;1842:1219–31.
- [15] LaFerla FM. Calcium dyshomeostasis and intracellular signalling in Alzheimer's disease. *Nat Rev Neurosci* 2002;3:862–72.
- [16] Smith IF, Green KN, LaFerla FM. Calcium dysregulation in Alzheimer's disease: recent advances gained from genetically modified animals. *Cell Calcium* 2005;38:427–37.
- [17] Berridge MJ. Calcium regulation of neural rhythms, memory and Alzheimer's disease. *J Physiol* 2014;592:281–93.
- [18] Supnet C, Bezprozvanny I. The dysregulation of intracellular calcium in Alzheimer disease. *Cell Calcium* 2010;47:183–9.
- [19] Stutzmann GE. The pathogenesis of Alzheimer's disease—is it a lifelong "Calciumopathy"? *Neuroscientist* 2007;13:546–59.
- [20] Bezprozvanny I, Mattson MP. Neuronal calcium mishandling and the pathogenesis of Alzheimer's disease. *Trends Neurosci* 2008;31:454–63.
- [21] Rizzuto R, De Stefani D, Raffaello A, Mammucari C. Mitochondria as sensors and regulators of calcium signalling. *Nat Rev Mol Cell Biol* 2012;13:566–78.
- [22] Rowland AA, Voeltz GK. Endoplasmic reticulum-mitochondria contacts: function of the junction. *Nat Rev Mol Cell Biol* 2012;13:607–25.
- [23] Zampese E, Fasolato C, Kipanyula MJ, Bortolozzi M, Pozzan T, Pizzo P. Presenilin 2 modulates endoplasmic reticulum (ER)-mitochondria interactions and Ca^{2+} cross-talk. *Proc Natl Acad Sci USA* 2011;108:2777–82.
- [24] Hedskog L, Pinho CM, Filadi R, et al. Modulation of the endoplasmic reticulum-mitochondria interface in Alzheimer's disease and related models. *Proc Natl Acad Sci USA* 2013;110:7916–21.
- [25] Filadi R, Greotti E, Turacchio G, Luini A, Pozzan T, Pizzo P. Mitofusin 2 ablation increases endoplasmic reticulum-mitochondria coupling. *Proc Natl Acad Sci USA* 2015;112:E2174–81.
- [26] Csordás G, Renken C, Várnai P, et al. Structural and functional features and significance of the physical linkage between ER and mitochondria. *J Cell Biol* 2006;174:915–21.
- [27] Qi H, Li L, Shuai J. Optimal microdomain crosstalk between endoplasmic reticulum and mitochondria for Ca^{2+} oscillations. *Sci Rep* 2015;5:7984.
- [28] Area-Gomez E, Castillo MdCL, Tambini MD, et al. Upregulated function of mitochondria-associated ER membranes in Alzheimer disease. *EMBO J* 2012;31:4106–23.
- [29] Schon EA, Area-Gomez E. Mitochondria-associated ER membranes in Alzheimer disease. *Mol Cell Neurosci* 2013;55:26–36.
- [30] Berridge MJ. Neuronal calcium signaling. *Neuron* 1998;21:13–26.

- [31] Bravo R, Vicencio JM, Parra V, et al. Increased ER-mitochondrial coupling promotes mitochondrial respiration and bioenergetics during early phases of ER stress. *J Cell Sci* 2011;124:2143–52.
- [32] Flood DG, Coleman PD. Neuron numbers and sizes in aging brain: comparisons of human, monkey, and rodent data. *Neurobiol Aging* 1988;9:453–63.
- [33] James BD, Glass TA, Caffo B, et al. Association of social engagement with brain volumes assessed by structural MRI. *J Aging Res* 2012;2012.
- [34] Martinelli C, Sartori P, Ledda M, Pannese E. A study of mitochondria in spinal ganglion neurons during life: quantitative changes from youth to extremely advanced age. *Tissue Cell* 2006;38:93–8.
- [35] Martinelli C, Sartori P, Ledda M, Pannese E. Mitochondria in perineuronal satellite cell sheaths of rabbit spinal ganglia: quantitative changes during life. *Cells Tissues Organs* 2007;186:141–6.
- [36] Morgan JE. Retina ganglion cell degeneration in glaucoma: an opportunity missed? A review. *Clin Exp Ophthalmol* 2012;40:364–8.
- [37] Wang X, Su B, Siedlak SL, et al. Amyloid- β overproduction causes abnormal mitochondrial dynamics via differential modulation of mitochondrial fission/fusion proteins. *Proc Natl Acad Sci USA* 2008;105:19318–23.
- [38] Cohen E, Dillin A. The insulin paradox: aging, proteotoxicity and neurodegeneration. *Nat Rev Neurosci* 2008;9:759–67.
- [39] Gleichmann M, Mattson MP. Neuronal calcium homeostasis and dysregulation. *Antioxid Redox Signal* 2011;14:1261–73.
- [40] Cheung K-H, Shineman D, Müller M, et al. Mechanism of Ca^{2+} disruption in Alzheimer's disease by presenilin regulation of InsP₃ receptor channel gating. *Neuron* 2008;58:871–83.
- [41] Cheung K-H, Mei L, Mak D-OD, et al. Gain-of-function enhancement of InsP₃ receptor modal gating by familial Alzheimer's disease-linked presenilin mutants in human cells and mouse neurons. *Sci Signal* 2010;3:ra22.
- [42] Stutzmann GE, Caccamo A, LaFerla FM, Parker I. Dysregulated IP₃ signaling in cortical neurons of knock-in mice expressing an Alzheimer's-linked mutation in presenilin1 results in exaggerated Ca^{2+} signals and altered membrane excitability. *J Neurosci* 2004;24:508–13.
- [43] Stutzmann GE, Smith I, Caccamo A, Oddo S, LaFerla FM, Parker I. Enhanced ryanodine receptor recruitment contributes to Ca^{2+} disruptions in young, adult, and aged Alzheimer's disease mice. *J Neurosci* 2006;26:5180–9.
- [44] Shilling D, Müller M, Takano H, et al. Suppression of InsP₃ receptor-mediated Ca^{2+} signaling alleviates mutant presenilin-linked familial Alzheimer's disease pathogenesis. *J Neurosci* 2014;34:6910–23.
- [45] Hung CH-L, Ho Y-S, Chang RC-C. Modulation of mitochondrial calcium as a pharmacological target for Alzheimer's disease. *Ageing Res Rev* 2010;9:447–56.
- [46] Wang C, Xie N, Wang Y, Li Y, Ge X, Wang M. Role of the mitochondrial calcium uniporter in rat hippocampal neuronal death after pilocarpine-induced status epilepticus. *Neurochem Res* 2015;40:1739–46.
- [47] Nakagawa T, Yuan J. Cross-talk between two cysteine protease families: activation of caspase-12 by calpain in apoptosis. *J Cell Biol* 2000;150:887–94.
- [48] Morishima N, Nakanishi K, Takenouchi H, Shibata T, Yasuhiko Y. An endoplasmic reticulum stress-specific caspase cascade in apoptosis cytochrome c-independent activation of caspase-9 by caspase-12. *J Biol Chem* 2002;277:34287–94.
- [49] Hitomi J, Katayama T, Taniguchi M, Honda A, Imaizumi K, Tohyama M. Apoptosis induced by endoplasmic reticulum stress depends on activation of caspase-3 via caspase-12. *Neurosci Lett* 2004;357:127–30.
- [50] Liu J, Liu MC, Wang K. Calpain in the CNS: from synaptic function to neurotoxicity. *Sci Signal* 2008;1:re1.
- [51] Bezprozvanny I. Calcium signaling and neurodegenerative diseases. *Trends Mol Med* 2009;15:89–100.
- [52] Norris CM, Kadish I, Blalock EM, et al. Calcineurin triggers reactive/inflammatory processes in astrocytes and is upregulated in aging and Alzheimer's models. *J Neurosci* 2005;25:4649–58.
- [53] Reese LC, Taglialatela G. A role for calcineurin in Alzheimer's disease. *Curr Neuropharmacol* 2011;9:685.
- [54] Storr SJ, Carragher NO, Frame MC, Parr T, Martin SG. The calpain system and cancer. *Nat Rev Cancer* 2011;11:364–74.
- [55] Wang H-G, Pathan N, Ethell IM, et al. Ca^{2+} -induced apoptosis through calcineurin dephosphorylation of BAD. *Science* 1999;284:339–43.
- [56] Youle RJ, Strasser A. The BCL-2 protein family: opposing activities that mediate cell death. *Nat Rev Mol Cell Biol* 2008;9:47–59.
- [57] Czabotar PE, Lessene G, Strasser A, Adams JM. Control of apoptosis by the BCL-2 protein family: implications for physiology and therapy. *Nat Rev Mol Cell Biol* 2014;15:49–63.
- [58] Galluzzi L, Blomgren K, Kroemer G. Mitochondrial membrane permeabilization in neuronal injury. *Nat Rev Neurosci* 2009;10:481–94.
- [59] Mattson MP, Magnus T. Ageing and neuronal vulnerability. *Nat Rev Neurosci* 2006;7:278–94.
- [60] Raynaud F, Marcilhac A. Implication of calpain in neuronal apoptosis. *FEBS J* 2006;273:3437–43.
- [61] Kitamura Y, Shimohama S, Kamoshima W, et al. Alteration of proteins regulating apoptosis, Bcl-2, Bcl-x, Bax, Bak, Bad, ICH-1 and CPP32, in Alzheimer's disease. *Brain Res* 1998;780:260–9.
- [62] D'Amelio M, Cavallucci V, Middei S, et al. Caspase-3 triggers early synaptic dysfunction in a mouse model of Alzheimer's disease. *Nat Neurosci* 2011;14:69–76.
- [63] Itkin A, Dupres V, Dufrêne YF, Bechinger B, Ruysschaert J-M, Raussens V. Calcium ions promote formation of amyloid β -peptide (1–40) oligomers causally implicated in neuronal toxicity of Alzheimer's disease. *PLoS ONE* 2011;6:e18250.
- [64] Kamer KJ, Mootha VK. The molecular era of the mitochondrial calcium uniporter. *Nat Rev Mol Cell Biol* 2015;16:545–53.
- [65] Cárdenas C, Miller RA, Smith I, et al. Essential regulation of cell bioenergetics by constitutive InsP₃ receptor Ca^{2+} transfer to mitochondria. *Cell* 2010;142:270–83.
- [66] Brookes PS, Yoon Y, Robotham JL, Anders M, Sheu S-S. Calcium, ATP, and ROS: a mitochondrial love-hate triangle. *Am J Physiol Cell Physiol* 2004;287:C817–33.
- [67] Alon U. Network motifs: theory and experimental approaches. *Nat Rev Genet* 2007;8:450–61.
- [68] Kinnally KW, Peixoto PM, Ryu S-Y, Dejean LM. Is mPTP the gatekeeper for necrosis, apoptosis, or both? *Biochim Biophys Acta* 2011;1813:616–22.
- [69] Rasola A, Bernardi P. Mitochondrial permeability transition in Ca^{2+} -dependent apoptosis and necrosis. *Cell Calcium* 2011;50:222–33.
- [70] Du H, Guo L, Fang F, et al. Cyclophilin D deficiency attenuates mitochondrial and neuronal perturbation and ameliorates learning and memory in Alzheimer's disease. *Nat Med* 2008;14:1097–105.
- [71] Nicotera P, Leist M, Manzo L. Neuronal cell death: a demise with different shapes. *Trends Pharmacol Sci* 1999;20:46–51.
- [72] Kanazawa I. How do neurons die in neurodegenerative diseases? *Trends Mol Med* 2001;7:339–44.
- [73] Müller M, Cárdenas C, Mei L, Cheung K-H, Foskett JK. Constitutive cAMP response element binding protein (CREB) activation by Alzheimer's disease presenilin-driven inositol triphosphate receptor (InsP₃R) Ca^{2+} signaling. *Proc Natl Acad Sci USA* 2011;108:13293–8.